

## Supplementary Appendix

This appendix has been provided by the authors to give readers additional information about their work.

Supplement to: Sherman KE, Flamm SL, Afdhal NH, et al. Response-guided telaprevir combination treatment for hepatitis C virus infection. N Engl J Med 2011;365:1014-24.

## APPENDIX

The members of the ILLUMINATE Study Group were as follows: **BELGIUM** – M. Adler, Erasme Hospital , Brussels; Jean Delwaide, Centro Hospitalo-Universitaire de Liege, Liege; Y. Horsmans, Clinique Universitaires Saint Luc, Brussels; H. van Vlierberghe, Ghent University Hospital, Ghent; **THE NETHERLANDS** – H.W. Reesink, Academic Medical Center of the University of Amsterdam, Amsterdam; C. Richter, Rijnstate Hospital, Arnhem; **AND THE UNITED STATES** – N. Afdhal, Beth Israel Deaconess Medical Center, Boston, MA; V. Araya, Albert Einstein Center for Liver Disease, Philadelphia, PA; S. Arora, University of New Mexico Division of Internal Medicine, Albuquerque, NM; L. Balart, Tulane University Health Sciences Center, New Orleans, LA; M. Bennett, Medical Associates Research Group, San Diego, CA; B. Berk, Dartmouth-Hitchcock Medical Center, Lebanon, NH; D. Bernstein, North Shore University Hospital, Manhasset, NY; J. Bloomer, University of Alabama at Birmingham, Birmingham, AL; R. Brown Jr., Columbia University Medical Center/NY Presbyterian New York, NY; N. Bzowej, CPMC Research Institute, San Francisco, CA; R. Chasen, Maryland Digestive Disease Research, Laurel, MD; J. Cochran, Birmingham Gastroenterology Associates, P.C. Birmingham, AL; J. Crippin, Washington University in St. Louis, St. Louis, MO; G. Davis, Liver Consultants of Texas, Dallas, TX; M. Davis, South Florida Center of Gastroenterology, Wellington, FL; E. DeJesus, Orlando Immunology, Orlando, FL; A. Di Bisceglie, St. Louis University, St. Louis, MO; D. Dieterich, Mount Sinai Medical Center, New York, NY; S. Esposito, Hepatobiliary Associates of New York, Bayside, NY; G. Everson, University of Colorado-Denver, Aurora, CO; S.L. Flamm, Northwestern University Medical Center, Chicago, IL; J. Franco, Medical College of Wisconsin, Milwaukee, WI; B. Freilich, Kansas City Gastroenterology & Hepatology, Kansas City, MO; M.W. Fried, University of North Carolina at Chapel Hill, Chapel Hill, NC, MWF is funded in part by NIH NIDDK Mid-Career Mentoring Award (K24 DK066144) and NIH NCRR 1UL1RR025747-01; R. Ghalib, Methodist Transplant Physicians, Dallas, TX; E. Godofsky, University Hepatitis Center at Bach & Godofsky, MD, PA, Sarasota, FL; S. Gordon, Henry Ford Columbus Center, Novi, MI; C. Howell, University of Maryland School of Medicine, Baltimore, MD; W. Hutson, University of Utah HSC, Salt Lake City, UT; I. Jacobson, Weill Cornell Medical College, New York, NY; M. Jonas, Consultants for Clinical Research, Cincinnati, OH; B. Kaufman, Atlantic Gastroenterology Associates, Egg Harbor Township, NJ; P. Kwo, Transplanatation CAQ Indiana University, Indianapolis, IN; E. Lawitz, Alamo Medical Research,

San Antonio, TX; M. Lucey, University of Wisconsin-Madison, Madison, WI; M. Lyons, Tacoma Digestive Disease Center, Tacoma, WA; P. Martin, University of Miami Division of Hepatology Center for Liver Disease, Miami, FL; A. Muir, Duke University Medical Center, Durham, NC; K. Mullen, MetroHealth Medical Center, Cleveland, OH; D. Nelson, University of Florida, Gainesville, FL; T. Nguyen, Research and Education, Inc. San Diego, CA; P. Pockros, Scripps Clinic, La Jolla, CA; F.F. Poordad, Cedars-Sinai Medical Center, Los Angeles, CA; R. Reindollar, Piedmont Healthcare, Statesville, NC; J. Reinus, Montefiore Medical Center, Bronx, NY; M. Rodriguez-Torres, San Juan, PR; L. Rossaro, University of California, Davis, Sacramento, CA; R. Rubin, Digestive Health Care GA, Atlanta, GA; R. Satyanarayana, Mayo Clinic Jacksonville, Jacksonville, FL; J. Scott, UW Harborview Medical Center, Seattle, WA; T. Sepe, University Gastroenterology, Providence, RI; N. Shah, Rush University Medical Center, Chicago, IL; A. Sheikh, GI Specialists of Georgia, PC, Marietta, GA; M. Sheikh, UCSF-Fresno, Fresno, CA; A. Sherker, MedStar Clinical Research, Washington, DC; K.E. Sherman, University of Cincinnati College of Medicine, Cincinnati, OH, efforts of KES supported in part by DK070528-06; J. Strohecker, Columbia Gastroenterology Associates, PA, Columbia, SC; M. Sulkowski, John Hopkins Rockland Physicians Practice & Research Group, Lutherville, MD; G. Szabo, UMASS Memorial Medical Center, Worcester, MA; N. Terrault, UCSF Medical Center, San Francisco, CA; H. Tobias, Concorde Medical Center, New York, NY; N. Tsai, Hawaii Medical Center, Honolulu, HI; H. Vargas, Mayo Hospital Clinic, Phoenix, AZ; K. Workowski, The Emory Clinic, Atlanta, GA; G. Wu, University of Connecticut Health Center, Farmington, CT; R. Yapp, Digestive Health Services, Downers Grove, IL; Z. Younes, GI Center of the MidSouth, Germantown, TN; Z. Younossi, Inova Health System, Falls Church, VA; A. Zaman, Oregon Health and Sciences University, Portland, OR.